Index of Authors

ABRAHAM, K. A. *see* ROGNES, A. 55–62
ACKERMANN, B. P. & AHLERS, J. Kinetics of alkaline phosphatase from pig kidney. Influence of complexing agents on stability and activity 151–157
AGOGBUA, S. I. O. & WYNN, C. H. Purification and properties of arylsulphatase B of human liver 415–421
AHLERS, J. *see* ACKERMANN, B. P. 151–157
AHLUWALIA, S. *see* HALLIWELL, B. 513–518
ALLEN, A. K. *see* SNARY, D. 75–78
ALLEN, M. B. *see* HOLROYDE, M. J. 363–373
ANDERSON, J. C. *see* LEE-OWN, V. 259–264
ANSTEE, D. J. *see* TANNER, M. J. A. 265–270, 271–277
ANTONINI, E. *see* COLOSIMO, A. 657–661
ATFIELD, G. *see* HOLNESS, N. J. 429–435, 437–446, 447–454

BAILEY, A. J. & SIMS, T. J. Chemistry of the collagen cross-links. Nature of the cross-links in the polymorphic forms of dermal collagen during development 211–215
BAILEY, G. S. & SHIPOLINI, R. A. Purification and properties of a kininogenin from the venom of *Vipera ammodytes ammodytes* 409–414
BARBER, M. J., BRAY, R. C., LOWE, D. J. & COUGHLAN, M. P. Studies by electron-paramagnetic-resonance spectroscopy and stopped-flow spectrophotometry on the mechanism of action of turkey liver xanthine dehydrogenase 297–307
BARDSLEY, W. G. The quantitative analysis of ligand binding and initial-rate data for allosteric and other complex enzyme mechanisms 101–117
BARNETT, W. A. *see* SPRAGG, S. P. 423–428
BEARDSLEE, R. A. *see* DONOVAN, J. W. 631–639
BECKER, R. R. *see* MCPARLAND, R. H. 597–606
BENTLEY, K. W. Quantitative determination of *N*-terminal amino acids of peptides and proteins with cobalt(III) chelates 137–138
BENTLEY, K. W. *see also* CHANG, J. Y. 607–611
BENZIMAN, M. *see* SWISSA, M. 173–179, 499–501
BLANCO, A., BURGOS, C., GEREZ DE BURGOS, N. M. & MONTAMAT, E. E. Properties of the testicular lactate dehydrogenase isoenzyme 165–172
BOULTER, D. *see* HASLETT, B. G. 33–38
BRAY, R. C. *see* BARBER, M. J. 297–307; DALTON, H. 287–295
BROWN, S. B. & KING, R. F. G. J. The structure of haem in pyridine/water mixtures and its implication in studies of haem catabolism 479–483
BROWN, S. B., SHILLCOCK, M. & JONES, P. Equilibrium and kinetic studies of the aggregation of porphyrins in aqueous solution 279–285
BROXMEYER, B. *see* ELCE, J. S. 223–232
BRUNORI, M. *see* COLOSIMO, A. 657–661
BRYCE, C. F. A., WILLIAMS, D. C., JOHN, R. A. & FASELLA, P. The anomalous kinetics of coupled aspartate aminotransferase and malate dehydrogenase. Evidence for compartmentation of oxaloacetate 571–577
BUONOCORE, V., POERIO, E., SILANO, V. & TOMASI, M. Physical and catalytic properties of α-amylase from *Tenebrio molitor* L. larvae 621–625

BURGOS, C. *see* BLANCO, A. 165–172
BURSTEIN, Y. *see* SCHECHTER, I. 543–550

CAREY, M. C., HIROM, P. C. & SMALL, D. M. A study of the physicochemical interactions between biliary lipids and chlorpromazine hydrochloride. Bile-salt precipitation as a mechanism of phenothiazine-induced bile secretory failure 519–531
CARREA, G. *see* LUGARO, G. 641–646
CASELLATO, M. M. *see* LUGARO, G. 641–646
CHANG, J. Y., CREASER, E. H. & BENTLEY, K. W. 4-*NN*-Dimethylaminoazobenzene 4′-isothiocyanate, a new chromophoric reagent for protein sequence analysis 607–611
CHANZY, H., FRANC, J.-M. & HERBAGE, D. High-angle electron diffraction of frozen hydrated collagen 139–140
CHESHER, J. M. E. *see* HOLROYDE, M. J. 351–361, 363–373
CHUDZIK, J. *see* DUBIN, A. 389–396; KOJ, A. 397–402
CLARKE, N. & DAWSON, R. M. C. Enzymic formation of glycerol 1:2-cyclic phosphate 745–747
CLEGG, R. A. Purification and some properties of nitrate reductase (EC 1.7.99.4) from *Escherichia coli* K12 533–541
COLOSIMO, A., BRUNORI, M. & ANTONINI, E. Electron transfer between soluble and immobilized mammalian cytochrome *c*. Equilibrium and kinetic studies on immobilized cytochrome *c* 657–661
COOREMAN, W. *see* SCHARPÉ, S. 505–507
CORNISH-BOWDEN, A. Estimation of the dissociation constants of enzyme–substrate complexes from steady-state measurements. Interpretation of pH-independence of K_m 455–461
CORNISH-BOWDEN, A. *see also* HOLROYDE, M. J. 363–373
CÔTÉ, R. H. & VALET, J.-P. Isolation, composition and reactivity of the neutral glycoproteins from human meconiums with specificities of the ABO and Lewis systems 63–73
COUGHLAN, M. P. *see* BARBER, M. J. 297–307
CREASER, E. H. *see* CHANG, J. Y. 607–611
CRUMPTON, M. J. *see* SNARY, D. 75–78

DALTON, H., LOWE, D. J., PAWLIK, R. T. & BRAY, R. C. Studies by electron-paramagnetic-resonance spectroscopy on the mechanism of action of xanthine dehydrogenase from *Veillonella alcalescens* 287–295
DARVEY, I. G. *see* WALKER, E. J. 329–337
DAVIES, B. H. *see* TAYLOR, R. F. 233–239
DAWSON, R. M. C. *see* CLARKE, N. 745–747
DAWSON, R. M. C. *see* HAZLEWOOD, G. P. 49–53
DELL, A. *see* MORRIS, H. R. 663–679
DICKENSON, C. J. & DICKINSON, F. M. Some properties of an alcohol dehydrogenase partially purified from baker's yeast grown without added zinc 309–319
DICKINSON, F. M. *see* DICKENSON, C. J. 309–319
DILS, R. *see* MANNING, R. 463–468
DIXON, H. B. F. The unreliability of estimates of group dissociation constants 627–629

DONOVAN, J. W., BEARDSLEE, R. A. & ROSS, K. D. Formation of monoferric ovotransferrins in the presence of chelates 631–639
DUBIN, A., KOJ, A. & CHUDZIK, J. Isolation and some molecular parameters of elastase-like neutral proteinases from horse blood leucocytes 389–396
DUBIN, A. *see also* KOJ, A. 397–402
DUNCAN, R. J. S., KLINE, J. E. & SOKOLOFF, L. Identity of brain alcohol dehydrogenase with liver alcohol dehydrogenase 561–566
DUNKER, A. K. & KENYON, A. J. Mobility of sodium dodecyl sulphate–protein complexes 191–197
DUNKERTON, J. & JAMES, S. P. Some observations on the $NADP^+$-linked oxidation of methylglyoxal catalysed by 2-oxoaldehyde dehydrogenase 503–504
DYKES, C. W. & KAY, J. Conversion of pepsinogen into pepsin is not a one-step process 141–144

EID, M. *see* SCHARPÉ, S. 505–507
ELCE, J. S. & BROXMEYER, B. γ-Glutamyltransferase of rat kidney. Simultaneous assay of the hydrolysis and transfer reactions with [*glutamate*-^{14}C]glutathione 223–232
ETHERINGTON, D. J. Bovine spleen cathepsin B1 and collagenolytic cathepsin. A comparative study of the properties of the two enzymes in the degradation of native collagen 199–209
EVANS, H. J. *see* MCPARLAND, R. H. 597–606
EVANSON, J. M. *see* WOOLLEY, D. E. 119–126

FASELLA, P. *see* BRYCE, C. F. A. 571–577
FAULKES, R. A. *see* SNARY, D. 75–78
FEINBERG, J. G. Modified methods of dialysis 721–723
FLANAGAN, T. *see* GREEN, N. 729–732
FOTHERGILL, L. A. & HODGSON, G. I. Yeast phosphoglycerate mutase. Cyanogen bromide cleavage and amino acid sequence of an active-site peptide 145–149
FRANC, J.-M. *see* CHANZY, H. 139–140

GEREZ DE BURGOS, N. M. *see* BLANCO, A. 165–172
GILES, I. G., POAT, P. C. & MUNDAY, K. A. Purification and properties of pyruvate kinase from the hepatopancreas of *Carcinus maenas* 127–134
GRAND, R. J. A. *see* SYSKA, H. 375–387
GREEN, G. *see* WOOLLEY, D. E. 119–126
GREEN, N. & FLANAGAN, T. The prediction of the conformation of membrane proteins from the sequence of amino acids 729–732
GREENWOOD, C. & MOORE, T. A. An investigation of protein conformation of cytochrome *c* by using cytochrome *c* insolubilized on to agarose gel 159–163
GRIVORYAN, N. A. *see* MARDANYAN, S. S. 725–727
GUEVARA, J. G. *see* MCPARLAND, R. H. 597–606

HALLIWELL, B. & AHLUWALIA, S. Hydroxylation of *p*-coumaric acid by horseradish peroxidase. The role of superoxide and hydroxyl radicals 513–518
HARRIS, J. I. *see* HILL, H. A. O. 551–560
HASLETT, B. G. & BOULTER, D. The *N*-terminal amino acid sequence of plastocyanin from *Stellaria media* L. An exercise to establish criteria for the identification of residues from a sequenator 33–38
HAZLEWOOD, G. P. & DAWSON, R. M. C. A phospholipid-deacylating system of bacteria active in a frozen medium 49–53
HERBAGE, D. *see* CHANZY, H. 139–140
HERZFELD, A. & RAPER, S. M. The heterogeneity of arginases in rat tissues 469–478
HILL, H. A. O., LOBB, R. R., SHARP, S. L., STOKES, A. M., HARRIS, J. I. & JACK, R. S. Metal-replacement studies in *Bacillus stearothermophilus* aldolase and a comparison of the mechanisms of class I and class II aldolases 551–560
HIROM, P. C. *see* CAREY, M. C. 519–531
HODGSON, G. I. *see* FOTHERGILL, L. A. 145–149
HOFMANN, T. *see* WANG, T.-T. 691–699; 701–712
HOLNESS, N. J. & ATFIELD, G. The extraction and purification of a cysteine transfer ribonucleic acid from baker's yeast 429–435
HOLNESS, N. J. & ATFIELD, G. The nucleotide sequence of cysteine transfer ribonucleic acid from baker's yeast. Identification of the products from partial degradation of the complete sequence 447–454
HOLNESS, N. J. & ATFIELD, G. The nucleotide sequence of cysteine transfer ribonucleic acid from baker's yeast. Products of complete digestion with pancreatic ribonuclease and ribonuclease T_1 437–446
HOLROYDE, M. J., ALLEN, M. B., STORER, A. C., WARSY, A. S., CHESHER, J. M. E., TRAYER, I. P., CORNISH-BOWDEN, A. & WALKER, D. G. The purification in high yield and characterization of rat hepatic glucokinase 363–373
HOLROYDE, M. J., CHESHER, J. M. E., TRAYER, I. P. & WALKER, D. G. Studies on the use of Sepharose–*N*-(6-aminohexanoyl)-2-amino-2-deoxy-D-glucopyranose for the large-scale purification of hepatic glucokinase 351–361
HORI, H. & TAMIYA, N. Preparation and activity of guanidinated or acetylated erabutoxins 217–222

IKAWA, S. & TAMMAR, A. R. Bile acids of sea snakes of the subfamily Viperinae and the biosynthesis of C-23-hydroxylated bile acids in liver homogenate fractions from the adder, *Vipera berus* (Linn.) 343–350
INGLEDEW, W. J. *see* OHNISHI, T. 39–48
ITO, S. & NICOL, J. A. C. Isolation of *S*-adenosyl-3-thiopropylamine from the eye of the sea catfish (*Arius felis*) 567–570
IUPAC COMMISSION ON THE NOMENCLATURE OF ORGANIC CHEMISTRY & IUPAC–IUB COMMISSION ON BIOCHEMICAL NOMENCLATURE. Nomenclature of cyclitols. Recommendations, 1973 23–31

JACK, R. S. *see* HILL, H. A. O. 551–560
JACOB, S. T., ROE, F. J. & ROSE, K. M. Chromatin-bound and free forms of poly(adenylic acid) polymerase in rat hepatic nuclei 733–735
JAMES, S. P. *see* DUNKERTON, J. 503–504
JOHN, R. A. *see* BRYCE, C. F. A. 571–577
JONES, M. N. & WILKINSON, A. The interaction between β-lactoglobulin and sodium *n*-dodecyl sulphate 713–718
JONES, P. *see* BROWN, S. B. 279–285

KATSANOS, N. A. *see* THOMOU, H. 241–247
KAY, J. *see* DYKES, C. W. 141–144
KENYON, A. J. *see* DUNKER, A. K. 191–197
KING, R. F. G. J. *see* BROWN, S. B. 479–483
KLINE, J. E. *see* DUNCAN, R. J. S. 561–566

KOJ, A., CHUDZIK, J. & DUBIN, A. Substrate specificity and modification of the active centre of elastase-like neutral proteinases from horse blood leucocytes 397–402
KOJ, A. *see also* DUBIN, A. 389–396

LAUWERS, A. *see* SCHARPÉ, S. 505–507
LEE-OWN, V. & ANDERSON, J. C. Interaction between proteoglycan subunit and type II collagen from bovine nasal cartilage, and the preferential binding of proteoglycan subunit to type I collagen 259–264
LÉONIS, J. *see* TRIBOUT, M. 89–91
LOBB, R. R. *see* HILL, H. A. O. 551–560
LOWE, D. J. *see* BARBER, M. J. 297–307; DALTON, H. 287–295
LUGARO, G., PASTA, P., CASELLATO, M. M., MAZZOLA, G. & CARREA, G. Chemical and physical properties of the human urinary glycoprotein with gastric antisecretory activity 641–646

MAEDA, N. & TAMIYA, N. Isolation, properties and amino acid sequences of three neurotoxins from the venom of a sea snake, *Aipysurus laevis* 79–87
MAGNUSSON, S. *see* MORRIS, H. R. 663–679
MALLIOPOULOU, T. B. *see* RAKITZIS, E. T. 737–739
MANNING, R., DILS, R. & MAYER, R. J. Purification and some properties of acetyl-coenzyme A carboxylase from rabbit mammary gland 463–468
MARDANYAN, S. S., GRIVORYAN, N. A. & NALBANDYAN, R. M. The effect of detergents on bovine adrenal ferredoxin 725–727
MARTIN, D. R. & WILLIAMS, R. J. P. Chemical nature and sequence of alamethicin 181–190
MAXWELL, I. H. Artifacts in the centrifugation of ribosomal and heterogeneous ribonucleic acid in '99%-dimethyl sulphoxide' gradients 509–512
MAYER, R. J. *see* MANNING, R. 463–468
MAZZOLA, G. *see* LUGARO, G. 641–646
MCDOUGALL, E. I. & STEWART, J. C. The whey proteins of the milk of red deer (*Cervus elaphus* L.). A homologue of bovine β-lactoglobulin 647–655
MCPARLAND, R. H., GUEVARA, J. G., BECKER, R. R. & EVANS, H. J. The purification and properties of the glutamine synthetase from the cytosol of soya-bean root nodules 597–606
MONTAMAT, E. E. *see* BLANCO, A. 165–172
MOORE, T. A. *see* GREENWOOD, C. 159–163
MORGANTHALER, J.-J. & PRICE, C. A. Density-gradient sedimentation in silica sols. Anomalous shifts in the banding densities of polystyrene 'latex' beads 487–490
MORRIS, H. R., DELL, A., PETERSEN, T. E., SOTTRUP-JENSEN, L. & MAGNUSSON, S. Mass-spectrometric identification and sequence location of the ten residues of the new amino acid (γ-carboxyglutamic acid) in the *N*-terminal region of prothrombin 663–679
MOSBY, L. J. *see* WOOD, E. J. 589–596
MUNDAY, K. A. *see* GILES, I. G. 127–134

NALBANDYAN, R. M. *see* MARDANYAN, S. S. 725–727
NICOL, J. A. C. *see* ITO, S. 567–570

OHNISHI, T., INGLEDEW, W. J. & SHIRAISHI, S. Resolution and functional characterization of two mitochondrial iron–sulphur centres of the 'high-potential iron–sulphur protein' type 39–48

PAREDES, S. *see* TRIBOUT, M. 89–91
PASTA, P. *see* LUGARO, G. 641–646
PATIL, N. B. β-Amylase-resistant amylose. Effect of urea on the limited hydrolysis of amylose by β-amylase 339–342
PAWLIK, R. T. *see* DALTON, H. 287–295
PERRY, S. V. *see* SYSKA, H. 375–387
PETERSEN, T. E. *see* MORRIS, H. R. 663–679
PHILLIPS, D. R. *see* POLYA, G. M. 681–690
PHILLIPS, N. C., ROBINSON, D. & WINCHESTER, B. G. Characterization of human liver α-D-mannosidase purified by affinity chromatography 579–587
PIETRUSZKO, R. & RYZEWSKI, C. N. A new subunit of horse liver alcohol dehydrogenase and subunit composition of the polymorphic form 249–257
POAT, P. C. *see* GILES, I. G. 127–134
POERIO, E. *see* BUONOCORE, V. 621–625
POLYA, G. M. & PHILLIPS, D. R. The occurrence in amino acid sequences of extensive informational symmetries based on possible codon–codon complementarity in the encoding polynucleotides 681–690
PRICE, C. A. *see* MORGANTHALER, J.-J. 487–490

RADHAKRISHNAN, A. N. *see* SIVAKAMI, S. 321–327
RAKITZIS, E. T. & MALLIOPOULOU, T. B. Inactivation of cathepsin D by dithiophosgene and by 2,2-dichloro-1,3-dithiacyclobutanone 737–739
RALSTON, G. B. *see* WALKER, E. J. 329–337
RAPER, S. M. *see* HERZFELD, A. 469–478
ROBINSON, D. *see* PHILLIPS, N. C. 579–587
ROBINSON, M. S. *see* WOOD, E. J. 589–596
ROCHE, J. J. *see* SPRAGG, S. P. 423–428
ROE, F. J. *see* JACOB, S. T. 733–735
ROGNES, A. & ABRAHAM, K. A. Substrate-binding ability of *Escherichia coli* ribonucleic acid polymerase in relation to its protein composition 55–62
ROSE, K. M. *see* JACOB, S. T. 733–735
ROSS, K. D. *see* DONOVAN, J. W. 631–639
RYZEWSKI, C. N. *see* PIETRUSZKO, R. 249–257

SAVOLAINEN, H. Isolation of the liver *N*-aspartyl-β-glucosaminidase in aspartylglucosaminuria 749–750
SCHARPÉ, S., EID, M., COOREMAN, W. & LAUWERS, A. α-1 Anti-trypsin, an inhibitor of renin 505–507
SCHECHTER, I. & BURSTEIN, Y. Identification of *N*-terminal methionine in the precursor of immunoglobulin light chain. Initiation of translation of messenger ribonucleic acid in plants and animals 543–550
SHARP, S. L. *see* HILL, H. A. O. 551–560
SHILLCOCK, M. *see* BROWN, S. B. 279–285
SHIPOLINI, R. A. *see* BAILEY, G. S. 409–414
SHIRAISHI, S. *see* OHNISHI, T. 39–48
SILANO, V. *see* BUONOCORE, V. 621–625
SIMS, T. J. *see* BAILEY, A. J. 211–215
SIVAKAMI, S. & RADHAKRISHNAN, A. N. Kinetic studies on glucoamylase of rabbit small intestine 321–327
SLABAS, A. R. & WALKER, D. A. Inhibition of spinach phosphoribulokinase by DL-glyceraldehyde 613–619
SMALL, D. M. *see* CAREY, M. C. 519–531
SMITH, P. I. & SWAN, G. A. A study of the supposed hydroxylation of tyrosine catalysed by peroxidase 403–408

SNARY, D., ALLEN, A. K., FAULKES, R. A. & CRUMPTON, M. J. Carbohydrate composition of lymphocyte plasma membrane from pig mesenteric lymph node 75–78
SOKOLOFF, L. *see* DUNCAN, R. J. S. 561–566
SOTTRUP-JENSEN, L. *see* MORRIS, H. R. 663–679
SPRAGG, S. P., WILCOX, J. K., ROCHE, J. J. & BARNETT, W. A. The association of yeast phosphoglycerate kinase (EC 2.7.2.3) 423–428
SPRING, T. G. *see* THOMAS, M. K. 741–744
STEWART, J. C. *see* MCDOUGALL, E. I. 647–655
STOKES, A. M. *see* HILL, H. A. O. 551–560
STORER, A. C. *see* HOLROYDE, M. J. 363–373
SÜDI, J. Chiral recognition of prochiral centres and general acid–base catalysis. Necessarily interrelated manifestations of active-site structure 491–493
SWAN, G. A. *see* SMITH, P. I. 403–408
SWISSA, M. & BENZIMAN, M. Factors affecting the activity of citrate synthase of *Acetobacter xylinum* and its possible regulatory role 173–179
SWISSA, M., WEINHOUSE, H. & BENZIMAN, M. Activities of citrate synthase and other enzymes of *Acetobacter xylinum in situ* and *in vitro* 499–501
SYSKA, H., WILKINSON, J. M., GRAND, R. J. A. & PERRY, S. V. The relationship between biological activity and primary structure of troponin I from white skeletal muscle of the rabbit 375–387

TAMIYA, N. *see* HORI, H. 217–222; MAEDA, N. 79–87
TAMMAR, A. R. *see* IKAWA, S. 343–350
TANNER, M. J. A. & ANSTEE, D. J. A method for the direct demonstration of the lectin-binding components of the human erythrocyte membrane 265–270
TANNER, M. J. A. & ANSTEE, D. J. The membrane change in En(a–) human erythrocytes. Absence of the major erythrocyte sialoglycoprotein 271–277
TAYLOR, R. F. & DAVIES, B. H. Triterpenoid carotenoids and related lipids. Triterpenoid carotenoid aldehydes from *Streptococcus faecium* UNH 564P 233–239
THOMAS, M. K. & SPRING, T. G. Inhibition of glycolytic enzymes by 2-phosphotartronate 741–744
THOMOU, H. & KATSANOS, N. A. The theory of the deoxyribonucleic acid–ribonucleic acid hybridization reaction 241–247
TOMASI, M. *see* BUONOCORE, V. 621–625
TRAYER, I. P. *see* HOLROYDE, M. J. 351–361, 363–373
TRIBOUT, M., PAREDES, S. & LÉONIS, J. Investigation of ribonuclease-catalysed kinetics by a micro-calorimetric method 89–91
TUCKER, J. S. *see* WOOLEY, D. E. 119–126

VALET, J.-P. *see* CÔTÉ, R. H. 63–73

WALKER, D. A. *see* SLABAS, A. R. 613–619
WALKER, D. G. *see* HOLROYDE, M. J. 351–361, 363–373
WALKER, E. J., RALSTON, G. B. & DARVEY, I. G. Further evidence for an allosteric model for ribonuclease 329–337
WANG, T.-T. & HOFMANN, T. Acyl and amino intermediates in reactions catalysed by pig pepsin. Analysis of transpeptidation products 691–699
WANG, T.-T. & HOFMANN, T. Effects of secondary binding by activator and inhibitor peptides on covalent intermediates of pig pepsin 701–712
WARSY, A. S. *see* HOLROYDE, M. J. 363–373
WEINHOUSE, H. *see* SWISSA, M. 499–501
WILCOX, J. K. *see* SPRAGG, S. P. 423–428
WILKINSON, A. *see* JONES, M. N. 713–718
WILKINSON, J. M. *see* SYSKA, H. 375–387
WILLIAMS, D. C. *see* BRYCE, C. F. A. 571–577
WILLIAMS, R. J. P. *see* MARTIN, D. R. 181–190
WILTON, D. C. Acrolein, an irreversible active-site-directed inhibitor of deoxyribose 5-phosphate aldolase? 495–497
WINCHESTER, B. G. *see* PHILLIPS, N. C. 579–587
WOOD, E. J., MOSBY, L. J. & ROBINSON, M. S. Characterization of the extracellular haemoglobin of *Haemopsis sanguisuga* (L.) 589–596
WOOLLEY, D. E., TUCKER, J. S., GREEN, G. & EVANSON, J. M. A neutral collagenase from human gastric mucosa 119–126
WOOSTER, M. S. & WRIGGLESWORTH, J. M. Adsorption of glyceraldehyde 3-phosphate dehydrogenase on condensed monolayers of phospholipid 93–100
WRIGGLESWORTH, J. M. *see* WOOSTER, M. S. 93–100
WYNN, C. H. *see* AGOGBUA, S. I. O. 415–421

Index of Subjects

Acetobacter xylinum, activities of citrate synthase and other enzymes in, *in situ* and *in vitro* (Swissa, M., Weinhouse, H. & Benziman, M.) 499–501

Acetobacter xylinum, factors affecting the activity of citrate synthase from, and its possible role in the regulation of the tricarboxylic acid cycle (Swissa, M. & Benziman, M.) 173–179

Acetylation, effects of guanidination and, on the activity of sea-snake (*Laticauda semifasciata*) venom erabutoxins (Hori, H. & Tamiya, N.) 217–222

Acetyl-coenzyme A carboxylase, purification and properties of, from rabbit mammary gland (Manning, R., Dils, R. & Mayer, R. J.) 463–468

Acid α-D-mannosidase, *see* α-D-Mannosidase, acid

Acrolein, irreversible inhibition by, of the activity of *Salmonella typhimurium* deoxyribose 5-phosphate aldolase (Wilton, D. C.) 495–497

Actin, binding sites of rabbit white-skeletal-muscle troponin I for troponin C and (Syska, H., Wilkinson, J. M., Grand, R. J. A. & Perry, S. V.) 375–387

Acyl intermediates, amino intermediates and, formed during transpeptidation catalysed by pig pepsin (Wang, T. T. & Hofmann, T.) 691–699

Acyl intermediates, effects of secondary binding by activator and inhibitor peptides on amino intermediates and, formed during transpeptidation catalysed by pig pepsin (Wang, T.-T. & Hofmann, T.) 701–712

Adder (*Vipera berus*), biosynthesis of C-23-hydroxylated bile acids by fractions of liver homogenate from (Ikawa, S. & Tammar, A. R.) 343–350

Adenosine triphosphate, effect of, and other factors affecting the activity of *Acetobacter xylinum* citrate synthase and its possible role in the regulation of the tricarboxylic acid cycle (Swissa, M. & Benziman, M.) 173–179

S-Adenosyl-3-thiopropylamine, isolation of, from the eye of the sea catfish (Ito, S. & Nicol, J. A. C.) 567–570

Adrenal gland, ox, effects of detergents on ferredoxin from (Mardanyan, S. S., Grigoryan, N. A. & Nalbandyan, R. M.) 725–727

Aipysurus laevis, *see* Sea snake

Alamethicin, chemical nature and amino acid sequence of (Martin, D. R. & Williams, R. J. P.) 181–190

Alcohol dehydrogenase, brain and liver, rat, evidence for the identity of (Duncan, R. J. S., Kline, J. E. & Sokoloff, L.) 561–566

Alcohol dehydrogenase, liver, horse, characterization of a new subunit of, and the subunit composition of the polymorphic form (Pietruszko, R. & Ryzewski, C. N.) 249–257

Alcohol dehydrogenase, partial purification and properties of, from baker's yeast grown without added zinc ions (Dickenson, C. J. & Dickinson, F. M.) 309–319

Aldolase, *see* Fructose diphosphate aldolase

Alkaline phosphatase, *see* Phosphatase, alkaline

Amino acid residues, criteria for the identification of, during determination of amino acid sequences of proteins with an automatic sequencer (Haslett, B. G. & Boulter, D.) 33–38

Amino acid residues, *N*-terminal, quantitative determination of, of peptides and proteins with cobalt(III) chelates 137–138

Amino acid sequence, chemical nature and, of alamethicin (Martin, D. R. & Williams, R. J. P.) 181–190

Amino acid sequence, cyanogen bromide cleavage and, of a peptide containing the active-site histidine residue of baker's-yeast phosphoglycerate mutase (Fothergill, L. A. & Hodgson, G. I.) 145–149

Amino acid sequence, *N*-terminal, mass-spectrometric identification and location of the ten γ-carboxyglutamic acid residues in, of ox prothrombin (Morris, H. R., Dell, A., Petersen, T. E., Sottrup-Jensen, L. & Magnusson, S.) 663–679

Amino acid sequence, *N*-terminal, of chickweed plastocyanin (Haslett, B. G. & Boulter, D.) 33–38

Amino acid sequences, criteria for the identification of amino acid residues during the determination of, of proteins with an automatic sequencer (Haslett, B. G. & Boulter, D.) 33–38

Amino acid sequences, isolation, properties and, of three neurotoxins from sea-snake (*Aipysurus laevis*) venom (Maeda, N. & Tamiya, N.) 79–87

Amino acid sequences of proteins, occurrence in, of extensive informational symmetries based on possible codon–codon complementarity in the encoding polynucleotides (Polya, G. M. & Phillips, D. R.) 681–690

Amino acid sequences of proteins, use of 4-*N*,*N*-dimethylaminoazobenzene 4′-isothiocyanate as a chromophoric reagent in the analysis of (Chang, J. Y., Creaser, C. H. & Bentley, K. W.) 607–611

Amino acid sequences, prediction of the conformation of membrane proteins from (Green, N. M. & Flanagan, M. T.) 729–732

Amino acids, possible role of rat kidney γ-glutamyltransferase in the transport of (Elce, J. S. & Broxmeyer, B.) 223–232

Amino intermediates, acyl intermediates and, formed during transpeptidation catalysed by pig pepsin (Wang, T. T. & Hofmann, T.) 691–699

Amino intermediates, effects of secondary binding by activator and inhibitor peptides on acyl intermediates and, formed during transpeptidation catalysed by pig pepsin (Wang, T.-T. & Hofmann, T.) 701–712

Ammonium ions, purification and properties of soya-bean root-nodule cytosol glutamine synthetase and its relevance to the assimilation of (McParland, R. H., Guevara, J. G., Becker, R. R. & Evans, H. J.) 597–606

α-Amylase, physical and chemical properties of, from the yellow mealworm (Buonocore, V., Poerio, E., Silano, V. & Tomasi, M.) 621–625

β-Amylase, sweet-potato-tuber, effect of urea on the limited hydrolysis of amylose by (Patil, N. B.) 339–342

Amylose, effect of urea on the limited hydrolysis of, by sweet-potato-tuber β-amylase (Patil, N. B.) 339–342

Anti-trypsin, α_1, plasma, human, characterization of, as an inhibitor of pig kidney renin (Scharpé, S., Eid, M., Cooreman, W. & Lauwers, A.) 505–507

Apoferredoxin, formation of, from ox adrenal-gland ferredoxin in the presence of detergents (Mardanyan, S. S., Grigoryan, N. A. & Nalbandyan, R. M.) 725–727

Arginases, heterogeneity of, from rat tissues (Herzfeld, A. & Raper, S. M.) 469–478

Arius felis, *see* Catfish, sea

Arylsulphatase B, purification and properties of, from human liver (Agogbua, S. I. O. & Wynn, C. H.) 415–421
Aspartate aminotransferase, heart, pig, evidence for the compartmentation of oxaloacetate from the anomalous kinetics of the coupled reaction catalysed by, and malate dehydrogenase (Bryce, C. F. A., Williams, D. C., John, R. A. & Fasella, P.) 571–577
N-Aspartyl-β-glucosaminidase, liver, human, isolation of, from a patient with aspartylglucosaminuria (Savolainen, H.) 749–750
Aspartylglucosaminuria, isolation of *N*-aspartyl-β-glucosaminidase from liver of a human patient with (Savolainen, H.) 749–750

Bacillus stearothermophilus, effects of the replacement of zinc by other metals in fructose diphosphate aldolase from, and a comparison of the mechanisms of action of class I and class II fructose diphosphate aldolases (Hill, H. A. O., Lobb, R. R., Sharp, S. L., Stokes, A. M., Harris, J. I. & Jack, R. S.) 551–560
Bean, soya (*Glycine max*), purification and properties of glutamine synthetase from the cytosol of root nodules of (McParland, R. H., Guevara, J. G., Becker, R. R. & Evans, H. J.) 597–606
Bile acids, C-23-hydroxylated, biosynthesis of, by fractions of liver homogenate from (Ikawa, S. & Tammar, A. R.) 343–350
Bile, human, physicochemical interactions between chlorpromazine hydrochloride and lipids of (Carey, M. C., Hirom, P. C. & Small, D. M.) 519–531
Bile salts, precipitation of, as a mechanism of phenothiazine-induced cholestasis in man (Carey, M. C., Hirom, P. C. & Small, D. M.) 519–531
Blood-group systems, ABO and Lewis, isolation, composition and reactivity of the neutral glycoproteins from human meconiums with specificities of (Côté, R. H. & Valet, J.-P.) 63–73
Brain, rat, evidence for the identity of rat liver alcohol dehydrogenase with the enzyme from (Duncan, R. J. S., Kline, J. E. & Sokoloff, L.) 561–566
Brush border, small-intestine, rabbit, purification and kinetic properties of glucoamylase/maltase complex from (Sivakami, S. & Radhakrishnan, A. N.) 321–327
Butyrivibrio sp., properties of a phospholipid-deacylating enzyme system in, that is active in a frozen medium (Hazlewood, G. P. & Dawson, R. M. C.) 49–53

Cadmium, effects of the replacement by, and other metals of zinc in *Bacillus stearothermophilus* fructose diphosphate aldolase and a comparison of the mechanisms of action of class I and class II fructose diphosphate aldolases (Hill, H. A. O., Lobb, R. R., Sharp, S. L., Stokes, A. M., Harris, J. I. & Jack, R. S.) 551–560
Caffeate, role of superoxide ion and hydroxyl radicals in the hydroxylation of *p*-coumarate to, by horseradish peroxidase (Halliwell, B. & Ahluwalia, S.) 513–518
Carbohydrate composition of pig mesenteric-lymph-node lymphocyte plasma membrane (Snary, D., Allen, A. K., Faulkes, R. A. & Crumpton, M. J.) 75–78
γ-Carboxyglutamic acid residues, ten, mass-spectrometric identification and location of, in the *N*-terminal amino acid sequence of ox prothrombin (Morris, H. R., Dell, A., Petersen, T. E., Sottrup-Jensen, L. & Magnusson, S.) 663–679
Carcinus maenas, *see* Crab
Carotenoids, triterpenoid, identification of 4,4′-diaponeurospen-4-al and 4,4′-diapolycopen-4-al as, in *Streptococcus faecium* UNH 564P (Taylor, R. F. & Davies, B. H.) 233–239
Cartilage, nasal-septum, ox, preferential binding of proteoglycan subunit to type I collagen rather than type II collagen from (Lee-Own, V. & Anderson, J. C.) 259–264
Catfish, sea (*Arius felis*), isolation of *S*-adenosyl-3-thiopropylamine from the eye of (Ito, S. & Nicol, J. A. C.) 567–570
Cathepsin B1, spleen, ox, comparative study of the properties of, and collagenolytic cathepsin in the degradation of native collagen (Etherington, D. J.) 199–209
Cathepsin, collagenolytic, spleen, ox, comparative study of the properties of, and cathepsin B1 in the degradation of native collagen (Etherington, D. J.) 199–209
Cathepsin D, spleen, ox, inactivation by dithiophosgene and 2,2-dichloro-1,3-dithiacyclobutanone of (Rakitzis, E. T. & Malliopoulou, T. B.) 737–739
Cells, hamster, polyoma-virus-transformed, artifacts in the centrifugation of ribosomal and heterogeneous ribonucleic acids from, in '99%-dimethyl sulphoxide' gradients (Maxwell, I. H.) 509–512
Cervus elaphus, *see* Deer, red
Chicken egg, formation of monoferric forms of ovotransferrin from, in the presence of iron-chelating agents (Donovan, J. W., Beardslee, R. A. & Ross, K. D.) 631–639
Chickweed (*Stellaria media*), *N*-terminal amino acid sequence of plastocyanin from (Haslett, B. G. & Boulter, D.) 33–38
Chloroplasts, spinach-leaf, inhibition by DL-glyceraldehyde of the activity of phosphoribulokinase of (Slabas, A. R. & Walker, D. A.) 613–619
Chlorpromazine hydrochloride, physicochemical interactions between human biliary lipids and (Carey, M. C. Hirom, P. C. & Small, D. M.) 519–531
Cholestasis, phenothiazine-induced, precipitation of bile salts as a mechanism of, in man (Carey, M. C., Hirom, P. C. & Small, D. M.) 519–531
Chromatin, forms of polyadenylate polymerase free in nuclear sap and bound to, in rat liver nuclei (Jacob, S. T., Roe, F. J. & Rose, K. M.) 733–735
Citrate synthase, *Acetobacter xylinum*, activities of, and other enzymes *in situ* and *in vitro* (Swissa, M., Weinhouse, H. & Benziman, M.) 499–501
Citrate synthase, *Acetobacter xylinum*, factors affecting the activity of, and its possible role in the regulation of the tricarboxylic acid cycle (Swissa, M. & Benziman, M.) 173–179
Citric acid cycle, *see* Tricarboxylic acid cycle
Cobalt(III) chelates, quantitative determination of *N*-terminal amino acid residues of peptides and proteins with (Bentley, K. W.) 137–138
Cobalt, effects of the replacement by, and other metals of zinc in *Bacillus stearothermophilus* fructose diphosphate aldolase and a comparison of the mechanisms of action of class I and class II fructose diphosphate aldolases (Hill, H. A. O., Lobb, R. R., Sharp, S. L., Stokes, A. M., Harris, J. I. & Jack, R. S.) 551–560

Codons, occurrence in amino acid sequences of proteins of extensive informational symmetries based on possible complementarity between, in the encoding polynucleotides (Polya, G. M. & Phillips, D. R.) 681–690
Collagen, cartilage, nasal-septum, ox, type I, preferential binding of proteoglycan subunit to, rather than the type II collagen (Lee-Own, V. & Anderson, J. C.) 259–264
Collagen, degradation of, by a neutral collagenase released by human gastric mucosa (Woolley, D. E., Tucker, J. S., Green, G. & Evanson, J. M.) 119–126
Collagen, hydrated, frozen, high-angle electron diffraction of (Chanzy, H., Franc, J.-M. & Herbage, D.) 139–140
Collagen, native, comparative study of the properties of ox spleen cathepsin B1 and collagenolytic cathepsin in the degradation of (Etherington, D. J.) 199–209
Collagen, skin, calf, changes in the nature of the cross-links in the polymorphic forms of, during foetal development (Bailey, A. J. & Sims, T. J.) 211–215
Collagenase, neutral, gastric-mucosa, human, properties of (Woolley, D. E., Tucker, J. S., Green, G. & Evanson, J. M.) 119–126
Concanavalin A, method for the direct demonstration of the human erythrocyte membrane components binding, and other lectins (Tanner, M. J. A. & Anstee, D. J.) 265–270
Coproporphyrin III, equilibrium and kinetic studies of the aggregation of, and other porphyrins in aqueous solution (Brown, S. B., Shillcock, M. & Jones, P.) 279–285
p-Coumarate, role of superoxide ion and hydroxyl radicals in the hydroxylation of, by horseradish peroxidase (Halliwell, B. & Ahluwalia, S.) 513–518
Crab (*Carcinus maenas*), purification and properties of pyruvate kinase from the hepatopancreas of (Giles, I. G., Poat, P. C. & Munday, K. A.) 127–134
Cyclic cytidine 2′:3′-monophosphate, *see* Cytidine 2′:3′-cyclic monophosphate
Cyclitols, nomenclature of (IUPAC Commission on the Nomenclature of Organic Chemistry & IUPAC–IUB Commission on Biochemical Nomenclature) 23–31
Cysteine transfer ribonucleic acid, baker's-yeast, derivation of the complete nucleotide sequence of (Holness, N. J. & Atfield, G.) 447–454
Cysteine transfer ribonucleic acid, baker's-yeast, nucleotide sequences of the products obtained by the complete digestion by pancreatic ribonuclease and ribonuclease T_1 of (Holness, N. J. & Atfield, G.) 437–446
Cysteine transfer ribonucleic acid, extraction and purification of, from baker's yeast (Holness, N. J. & Atfield, G.) 429–435
Cytidine 2′:3′-cyclic monophosphate, further evidence for an allosteric model for ox pancreas ribonuclease in the hydrolysis of (Walker, E. J., Ralston, G. B. & Darvey, I. G.) 329–337
Cytidine 2′:3′-cyclic monophosphate, use of a microcalorimetric method in an investigation of the kinetics of the hydrolysis of, by ribonuclease A (Tribout, M., Paredes, S. & Léonis, J.) 89–91
Cytochrome *c*, heart, horse, investigation of the protein conformation of cytochrome *c* by using, insolubilized on agarose gel (Greenwood, C. & Moore, T. A.) 159–163
Cytochrome *c*, heart, horse, soluble and immobilized, equilibrium and kinetic studies on electron transfer between (Colosimo, A., Brunori, M. & Antonini, E.) 657–661
Cytosol, root-nodule, soya-bean, purification and properties of glutamine synthetase from (McParland, R. H., Guevara, J. G., Becker, R. R. & Evans, H. J.) 597–606

Deer, red (*Cervus elaphus*), characterization of a homologue of cow β-lactoglobulin from the whey proteins of the milk of (McDougall, E. I. & Stewart, J. C.) 647–655
Deoxyribonucleic acid, theory of the hybridization reaction of ribonucleic acid and (Thomou, H. & Katsanos, N. A.) 241–247
Deoxyribose 5-phosphate aldolase, *Salmonella typhimurium*, irreversible inhibition by acrolein of the activity of (Wilton, D. C.) 495–497
Detergents, effects of, on ox adrenal-gland ferredoxin (Mardanyan, S. S., Grigoryan, N. A. & Nalbandyan, R. M.) 725–727
Deuteroporphyrin IX, equilibrium and kinetic studies of the aggregation of, and other porphyrins in aqueous solution (Brown, S. B., Shillcock, M. & Jones, P.) 279–285
Development, foetal, changes in the nature of the cross-links in the polymorphic forms of calf skin collagen during (Bailey, A. J. & Sims, T. J.) 211–215
Dialysis, modified methods of (Feinberg, J. G.) 721–723
4,4′-Diapolycopen-4-al, identification of 4,4′-diaponeurospen-4-al and, as triterpenoid carotenoids in *Streptococcus faecium* UNH 564P (Taylor, R. F. & Davies, B. H.) 233–239
4,4′-Diaponeurospen-4-al, identification of 4,4′-diapolycopen-4-al and, as triterpenoid carotenoids in *Streptococcus faecium* UNH 564P (Taylor, R. F. & Davies, B. H.) 233–239
2,2-Dichloro-1,3-dithiacyclobutanone, inactivation by dithiophosgene and, of ox spleen cathepsin D (Rakitzis, E. T. & Malliopoulou, T. B.) 737–739
Dihydroxyfumarate, effect of, on the conversion of tyrosine into 3,4-dihydroxyphenylalanine catalysed by horseradish peroxidase (Smith, P. I. & Swan, G. A.) 403–408
Dihydroxyfumarate, role of superoxide ion and hydroxyl radicals in the hydroxylation of *p*-coumarate to caffeate by horseradish peroxidase in the presence of (Halliwell, B. & Ahluwalia, S.) 513–518
3,4-Dihydroxyphenylalanine (dopa), study of the supposed conversion of tyrosine into, catalysed by horseradish peroxidase (Smith, P. I. & Swan, G. A.) 403–408
4-*NN*-Dimethylaminoazobenzene 4′-isothiocyanate, use of, as a chromophoric reagent in the analysis of amino acid sequences of proteins (Chang, J. Y., Creaser, E. H. & Bentley, K. W.) 607–611
Dissociation constants, group, unreliability of estimates of (Dixon, H. B. F.) 627–629
Dithiophosgene (2,2,4,4-tetrachloro-1,3-dithiacyclobutanone), inactivation by, and 2,2-dichloro-1,3-dithiacyclobutanone of ox spleen cathepsin D (Rakitzis, E. T. & Malliopoulou, T. B.) 737–739
Dodecyl sulphate, sodium, electrophoretic mobility of complexes of proteins and (Dunker, A. K. & Kenyon, A. J.) 191–197

Dodecyl sulphate, sodium, interaction between β-lactoglobulin and (Jones, M. N. & Wilkinson, A.) 713–718
Dopa (3,4-dihydroxyphenylalanine), study of the supposed conversion of tyrosine into, catalysed by horseradish peroxidase (Smith, P. I. & Swan, G. A.) 403–408

Egg, chicken, formation of monoferric forms of ovotransferrin from, in the presence of iron-chelating agents (Donovan, J. W., Beardslee, R. A. & Ross, K. D.) 631–639
Elastase, isolation and some molecular parameters of horse polymorphonuclear-leucocyte proteinases similar to (Dubin, A., Koj, A. & Chudzik, J.) 389–396
Elastase, substrate specificity and modification of the active centres of horse polymorphonuclear-leucocyte proteinases similar to (Koj, A., Chudzik, J. & Dubin, A.) 397–402
Electron diffraction, high-angle, of frozen hydrated collagen (Chanzy, H., Franc, J.-M. & Herbage, D.) 139–140
Electrons, equilibrium and kinetic studies on the transfer of, between soluble and immobilized horse heart cytochrome *c* (Colosimo, A., Brunori, M. & Antonini, E.) 657–661
Enolase, inhibition by 2-phosphotartronate of, and other glycolytic enzymes (Thomas, M. K. & Spring, T. G.) 741–744
Enzymes, chiral recognition of prochiral centres and general acid–base catalysis by (Südi, J.) 491–493
Enzymes, interpretation of the independence of pH of the Michaelis constants for reactions catalysed by (Cornish-Bowden, A.) 455–461
Enzymes, quantitative analysis of ligand binding and initial-rate data for allosteric and other complex mechanisms in reactions catalysed by (Bardsley, W. G.) 101–117
Erabutoxins, venom, sea-snake (*Laticauda semifasciata*), guanidinated and acetylated, preparation and activity of (Hori, H. & Tamiya, N.) 217–222
Erythrocruorin, extracellular, horseleech, characterization of (Wood, E. J., Mosby, L. J. & Robinson, M. S.) 589–596
Erythrocyte, human, En(a–), absence of the major sialoglycoprotein from the membrane of (Tanner, M. J. A. & Anstee, D. J.) 271–277
Erythrocytes, human, method for the direct demonstration of the lectin-binding components of the membrane of (Tanner, M. J. A. & Anstee, D. J.) 265–270
Escherichia coli K12, purification and properties of nitrate reductase from (Clegg, R. A.) 533–541
Escherichia coli M.R.E. 600, substrate-binding ability of ribonucleic acid polymerase from, in relation to its protein composition (Rognes, A. & Abraham, K. A.) 55–62
European sand viper (*Vipera ammodytes ammodytes*), purification and properties of kininogenin from the venom of (Bailey, G. S. & Shipolini, R. A.) 409–414
Eye, sea-catfish, isolation of *S*-adenosyl-3-thiopropylamine from (Ito, S. & Nicol, J. A. C.) 567–570

Ferredoxin, adrenal-gland, ox, effects of detergents on (Mardanyan, S. S., Grigoryan, N. A. & Nalbandyan, R. M.) 725–727
Fructose diphosphate aldolase, *Bacillus stearothermophilus*, effects of the replacement of zinc by other metals in, and a comparison of the mechanisms of action of class I and class II fructose diphosphate aldolases (Hill, H. A. O., Lobb, R. R., Sharp, S. L., Stokes, A. M., Harris, J. I. & Jack, R. S.) 551–560

Gastric mucosa, human, properties of a neutral collagenase released by (Woolley, D. E., Tucker, J. S., Green, G. & Evanson, J. M.) 119–126
Gastric secretion, chemical and physical properties of the human urinary glycoprotein inhibiting, in the rat (Lugaro, G , Pasta, P., Casellato, M. M., Mazzola, G. & Carrea, G.) 641–646
Glucoamylase/maltase complex, purification and kinetic properties of, from rabbit small intestine (Sivakami, S. & Radhakrishnan, A. N.) 321–327
Glucokinase, purification in high yield and characterization of, from rat liver (Holroyde, M. J., Allen, M. B., Storer, A. C., Warsy, A. S., Chesher, J. M. E., Trayer, I. P., Cornish-Bowden, A. & Walker, D. G.) 363–373
Glucokinase, studies on the use of Sepharose–*N*-(6-aminohexanoyl)-2-amino-2-deoxy-D-glucopyranose for the large-scale purification of, from rat liver (Holroyde, M. J., Chesher, J. M. E., Trayer, I. P. & Walker, D. G.) 351–361
Glutamine synthetase, purification and properties of, from soya-bean root-nodule cytosol (McParland, R. H., Guevara, J. G., Becker, R. R. & Evans, H. J.) 597–606
γ-Glutamyltransferase, kidney, rat, simultaneous assay of the hydrolysis and transfer reactions with glutathione catalysed by (Elce, J. S. & Broxmeyer, B.) 223–232
Glutathione, simultaneous assay of the hydrolysis and transfer reactions with, catalysed by rat kidney γ-glutamyltransferase (Elce, J. S. & Broxmeyer, B.) 223–232
DL-Glyceraldehyde, inhibition by, of the activity of spinach-leaf chloroplast phosphoribulokinase (Slabas, A. R. & Walker, D. A.) 613–619
Glyceraldehyde 3-phosphate dehydrogenase, skeletal-muscle, rabbit, adsorption of, on condensed monolayers of phospholipids (Wooster, M. S. & Wrigglesworth, J. M.) 93–100
Glycerokinase, *Acetobacter xylinum*, activities of citrate synthase and, and of other enzymes *in situ* and *in vitro* (Swissa, M., Weinhouse, H. & Benziman, M.) 499–501
Glycerol 1:2-cyclic phosphate, formation of, catalysed by rat kidney-cortex glycerophosphinicocholine diesterase (Clarke, N. & Dawson, R. M. C.) 745–747
Glycerophosphinicocholine diesterase, kidney-cortex, rat, formation of glycerol 1:2-cyclic phosphate catalysed by (Clarke, N. & Dawson, R. M. C.) 745–747
Glycine max, *see* Bean, soya
Glycolysis, inhibition by 2-phosphotartronate of enzymes involved in (Thomas, M. K. & Spring, T. G.) 741–744
Glycoprotein, major, absence of, from En(a–) human erythrocyte membrane (Tanner, M. J. A. & Anstee, D. J.) 271–277

Glycoprotein, urinary, human, chemical and physical properties of, with gastric anti-secretory activity (Lugaro, G., Pasta, P., Casellato, M. M., Mazzola, G. & Carrea, G.) 641–646
Glycoproteins, neutral, isolation, composition and reactivity of, from human meconiums with specificities of the ABO and Lewis blood-group systems (Côté, R. H. & Valet, J.-P.) 63–73
Group dissociation constants, unreliability of estimates of (Dixon, H. B. F.) 627–629
Guanidination, effects of acetylation and, on the activity of sea-snake (*Laticauda semifasciata*) venom erabutoxins (Hori, H. & Tamiya, N.) 217–222

Haem, structure of, in pyridine/water mixtures and its implication in studies of the catabolism of haem (Brown, S. B. & King, R. F. G. J.) 479–483
Haematoporphyrin IX, equilibrium and kinetic studies of the aggregation of, and other porphyrins in aqueous solution (Brown, S. B., Shillcock, M. & Jones, P.) 279–285
Haemoglobin, extracellular, horseleech, characterization of (Wood, E. J., Mosby, L. J. & Robinson, M. S.) 589–596
Haemopsis sanguisuga, *see* Horseleech
Heart, horse, equilibrium and kinetic studies on electron transfer between soluble and immobilized cytochrome *c* from (Colosimo, A., Brunori, M. & Antonini, E.) 657–661
Heart, horse, investigation of the protein conformation of cytochrome *c* by using cytochrome *c* from, insolubilized on agarose gel (Greenwood, C. & Moore, T. A.) 159–163
Heart, pig, evidence for the compartmentation of oxaloacetate from the anomalous kinetics of the coupled reaction catalysed by aspartate aminotransferase and malate dehydrogenase from (Bryce, C. F. A., Williams, D. C., John, R. A. & Fasella, P.) 571–577
Heart, pigeon and ox, resolution and functional characterization of two iron–sulphur centres of 'high-potential iron–sulphur protein' type in mitochondria from (Ohnishi, T., Ingledew, W. J. & Shiraishi, S.) 39–48
Hen, *see* Chicken
Hepatopancreas, crab (*Carcinus maenas*), purification and properties of pyruvate kinase from (Giles, I. G., Poat, P. C. & Munday, K. A.) 127–134
Heterodialysis, advantages and applications of (Feinberg, J. G.) 721–723
Histidine residue, active-site, cyanogen bromide cleavage and the amino acid sequence of a peptide containing, of baker's-yeast phosphoglycerate mutase (Fothergill, L. A. & Hodgson, G. I.) 145–149
Histidine residues, possible involvement of, in the active site of human liver arylsulphatase B (Agogbua, S. I. O. & Wynn, C. H.) 415–421
Horseleech (*Haemopsis sanguisuga*), characterization of the extracellular haemoglobin of (Wood, E. J., Mosby, L. J. & Robinson, M. S.) 589–596
Horseradish, role of superoxide ion and hydroxyl radicals in the hydroxylation of *p*-coumarate by peroxidase from (Halliwell, B. & Ahluwalia, S.) 513–518
cis-β-Hydroxyaquo(triethylenetetramine)cobalt(III) ion, quantitative determination of *N*-terminal amino acid residues of peptides and proteins with (Bentley, K. W.) 137–138
Hydroxyl radicals, role of superoxide ion radicals and, in the hydroxylation of *p*-coumarate by horseradish peroxidase (Halliwell, B. & Ahluwalia, S.) 513–518

Immunoglobulin light chain, myeloma MOPC-321, mouse, identification of an *N*-terminal methionine residue in the precursor of, biosynthesized by cell-free systems containing messenger ribonucleic acid from various plant and animal species (Schechter, I. & Burstein, Y.) 543–550
Instructions to authors, policy of the Journal and 1–21
Intestine, small, rabbit, purification and kinetic properties of glucoamylase/maltase complex from (Sivakami, S. & Radhakrishnan, A. N.) 321–327
Iodination, effects of, on the activities of elastase-like horse polymorphonuclear-leucocyte neutral proteinases (Koj, A., Chudzik, J. & Dubin, A.) 397–402
Ipomoea batatas, *see* Sweet potato
Iron-chelating agents, formation of monoferric forms of chicken ovotransferrin in the presence of (Donovan, J. W., Beardslee, R. A. & Ross, K. D.) 631–639
Iron–sulphur centres, mitochondrial, heart, pigeon and ox, two, resolution and functional characterization of, of 'high-potential iron–sulphur protein' type (Ohnishi, T., Ingledew, W. J. & Shiraishi, S.) 39–48
Iron–sulphur centres, studies by electron-paramagnetic-resonance spectroscopy and stopped-flow spectrophotometry of the role of molybdenum and, in the mechanism of action of turkey liver xanthine dehydrogenase (Barber, M. J., Bray, R. C., Lowe, D. J. & Coughlan, M. P.) 297–307
Iron–sulphur centres, studies by electron-paramagnetic-resonance spectroscopy of the role of molybdenum and, in the mechanism of action of *Veillonella alcalescens* xanthine dehydrogenase (Dalton, H., Lowe, D. J., Pawlik, R. T. & Bray, R. C.) 287–295
Iron–sulphur protein, high-potential, resolution and functional characterization of two pigeon and ox heart mitochondrial iron–sulphur centres typical of (Ohnishi, T., Ingledew, W. J. & Shiraishi, S.) 39–48
Isocyanates, inhibition by, of elastase-like horse polymorphonuclear-leucocyte neutral proteinases (Koj, A., Chudzik, J. & Dubin, A.) 397–402

Kallikrein, purification and properties of, from European-sand-viper venom (Bailey, G. S. & Shipolini, R. A.) 409–414
Kidney cortex, rat, formation of glycerol 1:2-cyclic phosphate catalysed by glycerophosphinicocholine diesterase from (Clarke, N. & Dawson, R. M. C.) 745–747
Kidney, pig, characterization of human plasma α_1 antitrypsin as an inhibitor of (Scharpé, S., Eid, M., Cooreman, W. & Lauwers, A.) 505–507
Kidney, pig, influence of metal ion-complexing agents on the stability and activity of alkaline phosphatase from (Ackermann, B. P. & Ahlers, J.) 151–157
Kidney, rat, heterogeneity of arginases from, and other tissues (Herzfeld, A. & Raper, S. M.) 469–478
Kidney, rat, simultaneous assay of the hydrolysis and transfer reactions with glutathione catalysed by γ-glutamyltransferase from (Elce, J. S. & Broxmeyer, B.) 223–232

Kinetics, quantitative analysis of ligand binding and initial-rate data for, for allosteric and other complex enzyme mechanisms (Bardsley, W. G.) 101–117
Kininogenin, purification and properties of, from European-sand-viper venom (Bailey, G. S. & Shipolini, R. A.) 409–414

Lactate dehydrogenase, chiral recognition of prochiral centres and general acid–base catalysis by enzymes exemplified by (Südi, J.) 491–493
Lactate dehydrogenase, testis, mouse, properties of the testicular isoenzyme from (Blanco, A., Burgos, C., Gerez de Burgos, N. M. & Montamat, E. E.) 165–172
β-Lactoglobulin, cow, characterization of a homologue of, from the whey proteins of red-deer milk (McDougall, E. I. & Stewart, J. C.) 647–655
β-Lactoglobulin, interaction between sodium dodecyl sulphate and (Jones, M. N. & Wilkinson, A.) 713–718
Laticauda semifasciata, *see* Sea snake
Leaves, spinach, inhibition by DL-glyceraldehyde of the activity of phosphoribulokinase of chloroplasts from (Slabas, A. R. & Walker, D. A.) 613–619
Lecithin, *see* Phosphatidylcholine
Lectins, method for the direct demonstration of the human erythrocyte membrane components binding (Tanner, M. J. A. & Anstee, D. J.) 265–270
Leech, horse-, *see* Horseleech
Leucocytes, polymorphonuclear, horse, isolation and some molecular parameters of elastase-like neutral proteinases from (Dubin, A., Koj, A. & Chudzik, J.) 389–396
Leucocytes, polymorphonuclear, substrate specificities and modification of the active centres of elastase-like neutral proteinases from (Koj, A., Chudzik, J. & Dubin, A.) 397–402
Ligands, quantitative analysis of the binding of, and initial-rate data for allosteric and other complex enzyme mechanisms (Bardsley, W. G.) 101–117
Ligands, unreliability of estimates of dissociation constants of groups involved in the binding of (Dixon, H. B. F.) 627–629
Light chain, immunoglobulin, myeloma MOPC-321, mouse, identification of an *N*-terminal methionine residue in the precursor of, biosynthesized by cell-free systems containing messenger ribonucleic acid from various plant and animal species (Schechter, I. & Burstein, Y.) 543–550
Lipids, biliary, human, physicochemical interactions between chlorpromazine hydrochloride and (Carey, M. C., Hirom, P. C. & Small, D. M.) 519–531
Liver, adder, biosynthesis of C-23-hydroxylated bile acids by fractions of homogenate of (Ikawa, S. & Tammar, A. R.) 343–350
Liver, horse, characterization of a new subunit of alcohol dehydrogenase from, and the subunit composition of the polymorphic form (Pietruszko, R. & Ryzewski, C. N.) 249–257
Liver, human, isolation of *N*-aspartyl-β-glucosaminidase from, of a patient with aspartylglucosaminuria (Savolainen, H.) 749–750
Liver, human, purification and properties of arylsulphatase B from (Agogbua, S. I. O. & Wynn, C. H.) 415–421
Liver, human, purification by affinity chromatography and characterization of acid α-D-mannosidase from (Phillips, N. C., Robinson, D. & Winchester, B. G.) 579–587
Liver, rat, chromatin-bound and free forms of polyadenylate polymerase in nuclei of (Jacob, S. T., Roe, F. J. & Rose, K. M.) 733–735
Liver, rat, evidence for the identity of rat brain alcohol dehydrogenase with the enzyme from (Duncan, R. J. S., Kline, J. E. & Sokoloff, L.) 561–566
Liver, rat, heterogeneity of arginases from, and other tissues (Herzfeld, A. & Raper, S. M.) 469–478
Liver, rat, purification in high yield and characterization of glucokinase from (Holroyde, M. J., Allen, M. B., Storer, A. C., Warsy, A. S., Chesher, J. M. E., Trayer, I. P., Cornish-Bowden, A. & Walker, D. G.) 363–373
Liver, rat, studies on the use of Sepharose–*N*-(6-aminohexanoyl)-2-amino-2-deoxy-D-glucopyranose for the large-scale purification of glucokinase from (Holroyde, M. J., Chesher, J. M. E., Trayer, I. P. & Walker, D. G.) 351–361
Liver, sheep, nicotinamide–adenine dinucleotide phosphate-linked oxidation of methylglyoxal catalysed by 2-oxo aldehyde dehydrogenase from (Dunkerton, J. & James, S. P.) 503–504
Liver, turkey, studies by electron-paramagnetic-resonance spectroscopy and stopped-flow spectrophotometry on the mechanism of action of xanthine dehydrogenase from (Barber, M. J., Bray, R. C., Lowe, D. J. & Coughlan, M. P.) 297–307
Lymph node, mesenteric, pig, carbohydrate composition of lymphocyte plasma membrane from (Snary, D., Allen, A. K., Faulkes, R. A. & Crumpton, M. J.) 75–78
Lymphocytes, mesenteric-lymph-node, pig, carbohydrate composition of plasma membrane from (Snary, D., Allen, A. K., Faulkes, R. A. & Crumpton, M. J.) 75–78
Lysozyme, electrophoretic mobility of complexes of sodium dodecyl sulphate and, and other proteins (Dunker, A. K. & Kenyon, A. J.) 191–197

Malate dehydrogenase, heart, pig, evidence for the compartmentation of oxaloacetate from the anomalous kinetics of the coupled reaction catalysed by pig heart aspartate aminotransferase and (Bryce, C. F. A., Williams, D. C., John, R. A. & Fasella, P.) 571–577
Maltase/glucoamylase complex, purification and kinetic properties of, from rabbit small intestine (Sivakami, S. & Radhakrishnan, A. N.) 321–327
Mammary gland, rabbit, purification and properties of acetyl-coenzyme A carboxylase from (Manning, R., Dils, R. & Mayer, R. J.) 463–468
Manganese, effects of the replacement by, and other metals of zinc in *Bacillus stearothermophilus* fructose diphosphate aldolase and a comparison of the mechanisms of action of class I and class II fructose diphosphate aldolases (Hill, H. A. O., Lobb, R. R., Sharp, S. L., Stokes, A. M., Harris, J. I. & Jack, R. S.) 551–560
α-D-Mannosidase, acid, purification by affinity chromatography and characterization of, from human liver (Phillips, N. C., Robinson, D. & Winchester, B. G.) 579–587

Mannosidosis, human, further evidence that deficiency of human liver acid α-D-mannosidase is a cause of (Phillips, N. C., Robinson, D. & Winchester, B. G.) 579–587
Mealworm, yellow (*Tenebrio molitor*), physical and catalytic properties of α-amylase from (Buonocore, V., Poerio, E., Silano, V. & Tomasi, M.) 621–625
Meconiums, human, isolation, composition and reactivity of the neutral glycoproteins from, with specificities of the ABO and Lewis blood-group systems (Côté, R. H. & Valet, J.-P.) 63–73
Melanogenesis, study on the role of peroxidase in (Smith, P. I. & Swan, G. A.) 403–408
Membrane, erythrocyte, human, En(a–), absence of the major sialoglycoprotein from (Tanner, M. J. A. & Anstee, D. J.) 271–277
Membrane, erythrocyte, human, method for the direct demonstration of the lectin-binding components of (Tanner, M. J. A. & Anstee, D. J.) 265–270
Membrane, *Escherichia coli* K12, purification and properties of nitrate reductase from (Clegg, R. A.) 533–541
Membrane, plasma, lymphocyte, mesenteric-lymph-node, pig, carbohydrate composition of (Snary, D., Allen, A. K., Faulkes, R. A. & Crumpton, M. J.) 75–78
Membranes, prediction of the conformation of proteins of, from their amino acid sequences (Green, N. M. & Flanagan, M. T.) 729–732
Messenger ribonucleic acid, *see* Ribonucleic acid, messenger
Metal ions, influence of complexing agents for, on the stability and activity of pig kidney alkaline phosphatase (Ackermann, B. P. & Ahlers, J.) 151–157
Methionine residue, *N*-terminal, identification of, in the precursor of mouse myeloma MOPC-321 immunoglobulin light chain biosynthesized by cell-free systems containing messenger ribonucleic acid from various plant and animal species (Schechter, I. & Burstein, Y.) 543–550
Methylglyoxal, nicotinamide–adenine dinucleotide phosphate-linked oxidation of, catalysed by sheep liver 2-oxo aldehyde dehydrogenase (Dunkerton, J. & James, S. P.) 503–504
Michaelis constants, interpretation of the independence of pH of, for enzyme-catalysed reactions (Cornish-Bowden, A.) 455–461
Milk, interaction between sodium dodecyl sulphate and β-lactoglobulin from (Jones, M. N. & Wilkinson, A.) 713–718
Milk, red-deer, characterization of a homologue of cow β-lactoglobulin from the whey proteins of (McDougall, E. I. & Stewart, J. C.) 647–655
Mitochondria, heart, pigeon and ox, resolution and functional characterization of two iron–sulphur centres of 'high-potential iron–sulphur protein' type in (Ohnishi, T., Ingledew, W. J. & Shiraishi, S.) 39–48
Molybdenum, studies by electron-paramagnetic-resonance spectroscopy and stopped-flow spectrophotometry of the role of iron–sulphur centres and, in the mechanism of action of turkey liver xanthine dehydrogenase (Barber, M. J., Bray, R. C., Lowe, D. J. & Coughlan, M. P.) 297–307
Molybdenum, studies by electron-paramagnetic-resonance spectroscopy of the role of iron–sulphur centres and, in the mechanism of action of *Veillonella alcalescens* xanthine dehydrogenase (Dalton, H., Lowe, D. J., Pawlik, R. T. & Bray, R. C.) 287–295
Mucosa, gastric, human, properties of a neutral collagenase released by (Woolley, D. E., Tucker, J. S., Green, G. & Evanson, J. M.) 119–126
Muscle, skeletal, rabbit, adsorption of glyceraldehyde 3-phosphate dehydrogenase from, on condensed monolayers of phospholipids (Wooster, M. S. & Wrigglesworth, J. M.) 93–100
Muscle, skeletal, rabbit, inhibition by 2-phosphotartronate of glycolytic enzymes from (Thomas, M. K. & Spring, T. G.) 741–744
Muscle, skeletal, white, rabbit, relationship between biological activity and primary structure of troponin I from (Syska, H., Wilkinson, J. M., Grand, R. J. A. & Perry, S. V.) 375–387
Myeloma MOPC-321, mouse, identification of an *N*-terminal methionine residue in the precursor of the immunoglobulin light chain of, biosynthesized by cell-free systems containing messenger ribonucleic acid from various plant and animal species (Schechter, I. & Burstein, Y.) 543–550

Nasal septum, ox, preferential binding of proteoglycan subunit to type I collagen rather than type II collagen from cartilage of (Lee-Own, V. & Anderson, J. C.) 259–264
Neurotoxins, three, isolation, properties and amino acid sequences of, from sea-snake (*Aipysurus laevis*) venom (Maeda, N. & Tamiya, N.) 79–87
Neurotoxins, venom, sea-snake (*Laticauda semifasciata*), guanidinated and acetylated, preparation and activity of (Hori, H. & Tamiya, N.) 217–222
Neutral proteinase, *see* Proteinase, neutral
Nitrate reductase, purification and properties of, from *Escherichia coli* K12 (Clegg, R. A.) 533–541
Nuclei, liver, rat, chromatin-bound and free forms of polyadenylate polymerase in (Jacob, S. T., Roe, F. J. & Rose, K. M.) 733–735
Nucleoside triphosphates, ability of *Escherichia coli* M.R.E. 600 ribonucleic acid polymerase to bind, in relation to its protein composition (Rognes, A. & Abraham, K. A.) 55–62
Nucleotide sequence of baker's-yeast cysteine transfer ribonucleic acid (Holness, N. J. & Atfield, G.) 447–454
Nucleotide sequences of the products obtained by the complete digestion by pancreatic ribonuclease and ribonuclease T_1 of baker's-yeast cysteine transfer ribonucleic acid (Holness, N. J. & Atfield, G.) 437–446

Ovotransferrin, chicken, formation of monoferric forms of, in the presence of iron-chelating agents (Donovan, J. W., Beardslee, R. A. & Ross, K. D.) 631–639
Oxaloacetate, evidence for the compartmentation of, from the anomalous kinetics of the coupled reaction catalysed by pig heart aspartate aminotransferase and malate dehydrogenase (Bryce, C. F. A., Williams, D. C., John, R. A. & Fasella, P.) 571–577

2-Oxo aldehyde dehydrogenase, liver, sheep, nicotinamide–adenine dinucleotide phosphate-linked oxidation of methylglyoxal catalysed by (Dunkerton, J. & James, S. P.) 503–504

Pancreas, ox, further evidence for an allosteric model for ribonuclease from, in the hydrolysis of cytidine 2′:3′-cyclic monophosphate (Walker, E. J., Ralston, G. B. & Darvey, I. G.) 329–337
Pepsin, pig, acyl and amino intermediates formed during transpeptidation catalysed by (Wang, T. T. & Hofmann, T.) 691–699
Pepsin, pig, effects of secondary binding by activator and inhibitor peptides on covalent intermediates formed during transpeptidation catalysed by (Wang, T.-T. & Hofmann, T.) 701–712
Pepsin, pig, evidence that the conversion of pepsinogen into, is not a one-step process (Dykes, C. W. & Kay, J.) 141–144
Pepsinogen, pig, evidence that the conversion of, into pepsin is not a one-step process (Dykes, C. W. & Kay, J.) 141–144
Pepstatin, use of, to obtain evidence that the conversion of pig pepsinogen into pepsin is not a one-step process (Dykes, C. W. & Kay, J.) 141–144
Peptides, acyl and amino intermediates formed during transpeptidation of, catalysed by pig pepsin (Wang, T. T. & Hofmann, T.) 691–699
Peptides, effects of secondary binding by activator and inhibitor peptides on covalent intermediates formed during transpeptidation of, catalysed by pig pepsin (Wang, T.-T. & Hofmann, T.) 701–712
Peptides, quantitative determination of *N*-terminal amino acid residues of proteins and, with cobalt(III) chelates (Bentley, K. W.) 137–138
Peroxidase, horseradish, role of superoxide ion and hydroxyl radicals in the hydroxylation of *p*-coumarate by (Halliwell, B. & Ahluwalia, S.) 513–518
Peroxidase, horseradish, study of the supposed conversion of tyrosine into 3,4-dihydroxyphenylalanine catalysed by (Smith, P. I. & Swan, G. A.) 403–408
pH, interpretation of the independence of, of the Michaelis constants for enzyme-catalysed reactions (Cornish-Bowden, A.) 455–461
Phenothiazines, precipitation of bile salts as a mechanism of cholestasis induced by, in man (Carey, M. C., Hirom, P. C. & Small, D. M.) 519–531
Phenylalaninol residue, side-chain, identification of, in alamethicin (Martin, D. R. & Williams, R. J. P.) 181–190
Phosphatase, alkaline, kidney, pig, influence of metal ion-complexing agents on the stability and activity of (Ackermann, B. P. & Ahlers, J.) 151–157
Phosphatidic acid, adsorption of rabbit skeletal-muscle glyceraldehyde 3-phosphate dehydrogenase on condensed monolayers of (Wooster, M. S. & Wrigglesworth, J. M.) 93–100
Phosphatidylcholine, adsorption of rabbit skeletal-muscle glyceraldehyde 3-phosphate dehydrogenase on condensed monolayers of phosphatidic acid in the presence of (Wooster, M. S. & Wrigglesworth, J. M.) 93–100
Phosphatidylcholine, decomposition of, by a *Butyrivibrio* sp. phospholipid-deacylating enzyme system that is active in a frozen medium (Hazlewood, G. P. & Dawson, R. M. C.) 49–53
Phosphoglycerate kinase, yeast, association of (Spragg, S. P., Wilcox, J. K., Roche, J. J. & Barnett, W. A.) 423–428
Phosphoglycerate mutase, baker's-yeast, cyanogen bromide cleavage and the amino acid sequence of a peptide containing the active-site histidine residue of (Fothergill, L. A. & Hodgson, G. I.) 145–149
Phosphoglycerate mutase, inhibition by 2-phosphotartronate of, and other glycolytic enzymes (Thomas, M. K. & Spring, T. G.) 741–744
Phospholipase, *Butyrivibrio* sp., properties of, that is active in a frozen medium (Hazlewood, G. P. & Dawson, R. M. C.) 49–53
Phospholipids, adsorption of rabbit skeletal-muscle glyceraldehyde 3-phosphate dehydrogenase on condensed monolayers of (Wooster, M. S. & Wrigglesworth, J. M.) 93–100
Phospholipids, biliary, human, physicochemical interactions between chlorpromazine hydrochloride and (Carey, M. C., Hirom, P. C. & Small, D. M.) 519–531
Phospholipids, properties of a *Butyrivibrio* sp. enzyme system deacylating, that is active in a frozen medium (Hazlewood, G. P. & Dawson, R. M. C.) 49–53
Phosphoribulokinase, chloroplast, spinach-leaf, inhibition by DL-glyceraldehyde of the activity of (Slabas, A. R. & Walker, D. A.) 613–619
2-Phosphotartronate, inhibition by, of glycolytic enzymes (Thomas, M. K. & Spring, T. G.) 741–744
Photosynthesis, investigation of the sites of the inhibition by DL-glyceraldehyde of, by spinach-leaf chloroplasts (Slabas, A. R. & Walker, D. A.) 613–619
Phytohaemagglutinin, method for the direct demonstration of the human erythrocyte membrane components binding, and other lectins (Tanner, M. J. A. & Anstee, D. J.) 265–270
Plasma, human, characterization of α_1 anti-trypsin from, as an inhibitor of pig kidney renin (Scharpé, S., Eid, M., Cooreman, W. & Lauwers, A.) 505–507
Plasma membrane, *see* Membrane, plasma
Plasmacytoma MOPC-321, mouse, identification of an *N*-terminal methionine residue in the precursor of the immunoglobulin light chain of, biosynthesized by cell-free systems containing messenger ribonucleic acid from various plant and animal species (Schechter, I. & Burstein, Y.) 543–550
Plastocyanin, chickweed, *N*-terminal amino acid sequence of (Haslett, B. G. & Boulter, D.) 33–38
Policy of the Journal and instructions to authors 1–21
Polyadenylate polymerase, nuclear, liver, rat, chromatin-bound and free forms of (Jacob, S. T., Roe, F. J. & Rose, K. M.) 733–735
Polymorphonuclear leucocytes, *see* Leucocytes, polymorphonuclear
Polynucleotides, encoding, occurrence in amino acid sequences of proteins of extensive informational symmetries based on possible codon–codon complementarity in (Polya, G. M. & Phillips, D. R.) 681–690

Polysaccharides, physical and chemical properties of α-amylase from the yellow mealworm and their relevance to the digestion of, by the insect larva (Buonocore, V., Poerio, E., Silano, V. & Tomasi, M.) 621–625

Polystyrene beads, anomalous shifts in the banding densities of, during density-gradient sedimentation in silica sols (Morgenthaler, J.-J. & Price, C. A.) 487–490

Porphyrins, equilibrium and kinetic studies of the aggregation of, in aqueous solution (Brown, S. B., Shillcock, M. & Jones, P.) 279–285

Pregnancy, chemical and physical properties of the glycoprotein with gastric anti-secretory activity obtained from human urine during (Lugaro, G., Pasta, P., Casellato, M. M., Mazzola, G. & Carrea, G.) 641–646

Protease, *see* Proteinase

Protein, composition of, of *Escherichia coli* M.R.E. 600 ribonucleic acid polymerase and its relation to the substrate-binding ability of the enzyme (Rognes, A. & Abraham, K. A.) 55–62

Protein, iron–sulphur, high-potential, resolution and functional characterization of two pigeon and ox heart mitochondrial iron–sulphur centres typical of (Ohnishi, T., Ingledew, W. J. & Shiraishi, S.) 39–48

Protein (β-lactoglobulin), interaction between sodium dodecyl sulphate and (Jones, M. N. & Wilkinson, A.) 713–718

Proteinases, neutral, elastase-like, isolation and some molecular parameters of, from horse polymorphonuclear leucocytes (Dubin, A., Koj, A. & Chudzik, J.) 389–396

Proteinases, neutral, elastase-like, substrate specificity and modification of the active centres of, from horse polymorphonuclear leucocytes (Koj, A., Chudzik, J. & Dubin, A.) 397–402

Protein–polysaccharide, *see* Proteoglycan

Proteins, criteria for the identification of amino acid residues during the determination of amino acid sequences of, with an automatic sequencer (Haslett, B. G. & Boulter, D.) 33–38

Proteins, electrophoretic mobility of complexes of sodium dodecyl sulphate and (Dunker, A. K. & Kenyon, A. J.) 191–197

Proteins, membrane, prediction of the conformation of, from their amino acid sequences (Green, N. M. & Flanagan, M. T.) 729–732

Proteins, occurrence in amino acid sequences of, of extensive informational symmetries based on possible codon–codon complementarity in the encoding polynucleotides (Polya, G. M. & Phillips, D. R.) 681–690

Proteins, quantitative determination of *N*-terminal amino acid residues of peptides and, with cobalt(III) chelates (Bentley, K. W.) 137–138

Proteins, simultaneous fractionation and concentration of, by heterodialysis (Feinberg, J. G.) 721–723

Proteins, use of 4-*NN*-dimethylaminoazobenzene 4′-isothiocyanate as a chromophoric reagent in the analysis of the amino acid sequences of (Chang, J. Y., Creaser, E. H. & Bentley, K. W.) 607–611

Proteins, whey, milk, red-deer, characterization of a homologue of cow β-lactoglobulin from (McDougall, E. I. & Stewart, J. C.) 647–655

Proteoglycan subunit, preferential binding of, to type I collagen rather than type II collagen from ox nasal-septum cartilage (Lee-Own, V. & Anderson, J. C.) 259–264

Prothrombin, ox, mass-spectrometric identification and location of the ten γ-carboxyglutamic acid residues in the *N*-terminal amino acid sequence of (Morris, H. R., Dell, A., Petersen, T. E., Sottrup-Jensen, L. & Magnusson, S.) 663–679

Protoporphyrin IX, equilibrium and kinetic studies of the aggregation of, and other porphyrins in aqueous solution (Brown, S. B., Shillcock, M. & Jones, P.) 279–285

Pyruvate kinase, inhibition by 2-phosphotartronate of, and other glycolytic enzymes (Thomas, M. K. & Spring, T. G.) 741–744

Pyruvate kinase, purification and properties of, from crab (*Carcinus maenas*) hepatopancreas (Giles, I. G., Poat, P. C. & Munday, K. A.) 127–134

Red blood cells, *see* Erythrocytes

Red deer, *see* Deer, red

Renin, kidney, pig, characterization of human plasma α_1 anti-trypsin as an inhibitor of (Scharpé, S., Eid, M., Cooreman, W. & Lauwers, A.) 505–507

Respiratory chain, mitochondrial, heart, pigeon and ox, resolution and functional characterization of two iron–sulphur centres of 'high-potential iron–sulphur protein' type in (Ohnishi, T., Ingledew, W. J. & Shiraishi, S.) 39–48

Ribonuclease A, use of a micro calorimetric method in an investigation of the kinetics of the reaction catalysed by (Tribout, M., Paredes, S. & Léonis, J.) 89–91

Ribonuclease, pancreas, ox, further evidence for an allosteric model for, in the hydrolysis of cytidine 2′:3′-cyclic monophosphate (Walker, E. J., Ralston, G. B. & Darvey, I. G.) 329–337

Ribonucleic acid, messenger, identification of an *N*-terminal methionine residue in the precursor of mouse myeloma MOPC-321 immunoglobulin light chain biosynthesized by cell-free systems containing, from various plant and animal species (Schechter, I. & Burstein, Y.) 543–550

Ribonucleic acid polymerase, *Escherichia coli* M.R.E. 600, substrate-binding ability of, in relation to its protein composition (Rognes, A. & Abraham, K. A.) 55–62

Ribonucleic acid, theory of the hybridization reaction of deoxyribonucleic acid and (Thomou, H. & Katsanos, N. A.) 241–247

Ribonucleic acid, transfer, cysteine, baker's-yeast, derivation of the complete nucleotide sequence of (Holness, N. J. & Atfield, G.) 447–454

Ribonucleic acid, transfer, cysteine, baker's-yeast, nucleotide sequences of the products obtained by the complete digestion by pancreatic ribonuclease and ribonuclease T_1 of (Holness, N. J. & Atfield, G.) 437–446

Ribonucleic acid, transfer, cysteine, extraction and purification of, from baker's yeast (Holness, N. J. & Atfield, G.) 429–435

Ribonucleic acid, yeast, use of a micro calorimetric method in an investigation of the kinetics of the hydrolysis of, catalysed by ribonuclease A (Tribout, M., Paredes, S. & Léonis, J.) 89–91
Ribonucleic acids, messenger, encoding, occurrence in amino acid sequences of proteins of extensive informational symmetries based on possible codon–codon complementary in (Polya, G. M. & Phillips, D. R.) 681–690
Ribonucleic acids, ribosomal and heterogeneous, artifacts in the centrifugation of, in '99%-dimethyl sulphoxide' gradients (Maxwell, I. H.) 509–512
Ribosomal ribonucleic acid, *see* Ribonucleic acid, ribosomal
Ricin, method for the direct demonstration of the human erythrocyte membrane components binding, and other lectins (Tanner, M. J. A. & Anstee, D. J.) 265–270
Root nodules, soya-bean, purification and properties of glutamine synthetase from the cytosol of (McParland, R. H., Guevara, J. G., Becker, R. R. & Evans, H. J.) 597–606

Saccharomyces cerevisiae, cyanogen bromide cleavage and the amino acid sequence of a peptide containing the active-site histidine residue of phosphoglycerate mutase from (Fothergill, L. A. & Hodgson, G. I.) 145–149
Saccharomyces cerevisiae, derivation of the complete nucleotide sequence of cysteine transfer ribonucleic acid from (Holness, N. J. & Atfield, G.) 447–454
Saccharomyces cerevisiae, extraction and purification of a cysteine transfer ribonucleic acid from (Holness, N. J. & Atfield, G.) 429–435
Saccharomyces cerevisiae, nucleotide sequences of the products obtained by the complete digestion by pancreatic ribonuclease and ribonuclease T_1 of cysteine transfer ribonucleic acid from (Holness, N. J. & Atfield, G.) 437–446
Saccharomyces cerevisiae, partial purification and properties of alcohol dehydrogenase from, grown without added zinc ions (Dickenson, C. J. & Dickinson, F. M.) 309–319
Salivary gland, submaxillary, rat, heterogeneity of arginases from, and other tissues (Herzfeld, A. & Raper, S. M.) 469–478
Salmonella typhimurium, irreversible inhibition by acrolein of the activity of deoxyribose 5-phosphate aldolase from (Wilton, D. C.) 495–497
Sea catfish, *see* Catfish, sea
Sea snake (*Aipysurus laevis*), isolation, properties and amino acid sequences of three neurotoxins from the venom of (Maeda, N. & Tamiya, N.) 79–87
Sea snake (*Laticauda semifasciata*), preparation and activity of guanidinated and acetylated erabutoxins from the venom of (Hori, H. & Tamiya, N.) 217–222
Sialoglycoprotein, major, absence of, from En(a–) human erythrocyte membrane (Tanner, M. J. A. & Anstee, D. J.) 271–277
Sialoglycoproteins, isolation, composition and reactivity of, from human meconiums with specificities of the ABO and Lewis blood-group systems (Côté, R. H. & Valet, J.-P.) 63–73
Silica sols, anomalous shifts in the banding densities of polystyrene beads during density-gradient sedimentation in (Morgenthaler, J.-J. & Price, C. A.) 487–490
Skeletal muscle, *see* Muscle, skeletal
Skin, calf, changes in the nature of the cross-links in the polymorphic forms of collagen from, during foetal development (Bailey, A. J. & Sims, T. J.) 211–215
Small intestine, *see* Intestine, small
Sodium dodecyl sulphate, electrophoretic mobility of complexes of proteins and (Dunker, A. K. & Kenyon, A. J.) 191–197
Sodium dodecyl sulphate, interaction between β-lactoglobulin and (Jones, M. N. & Wilkinson, A.) 713–718
Soya bean, *see* Bean, soya
Spinach (*Spinacia oleracea*) leaves, inhibition by DL-glyceraldehyde of the activity of phosphoribulokinase of chloroplasts from (Slabas, A. R. & Walker, D. A.) 613–619
Spinacia oleracea, *see* Spinach
Spleen, ox, comparative study of the properties of cathepsin B1 and collagenolytic cathepsin from, in the degradation of native collagen (Etherington, D. J.) 199–209
Spleen, ox, inactivation by dithiophosgene and 2,2-dichloro-1,3-dithiacyclobutanone of cathepsin D from (Rakitzis, E. T. & Malliopoulou, T. B.) 737–739
Stellaria media, *see* Chickweed
Stomach, human, properties of a neutral collagenase released by mucosa of (Woolley, D. E., Tucker, J. S., Green, G. & Evanson, J. M.) 119–126
Stomach, rat, chemical and physical properties of the human urinary glycoprotein inhibiting gastric secretion by (Lugaro, G., Pasta, P., Casellato, M. M., Mazzola, G. & Carrea, G.) 641–646
Streptococcus faecium UNH 564P, identification of 4,4′-diaponeurospen-4-al and 4,4′-diapolycopen-4-al as triterpenoid carotenoids in (Taylor, R. F. & Davies, B. H.) 233–239
Submaxillary gland, rat, heterogeneity of arginases from, and other tissues (Herzfeld, A. & Raper, S. M.) 469–478
Supernatant fraction, *see* Cytosol
Superoxide ion radicals, role of hydroxyl radicals and, in the hydroxylation of *p*-coumarate by horseradish peroxidase (Halliwell, B. & Ahluwalia, S.) 513–518
Sweet-potato (*Ipomoea batatas*) tubers, effect of urea on the limited hydrolysis of amylose by β-amylase from (Patil, N. B.) 339–342

Temperature, low, properties of a *Butyrivibrio* sp. phospholipid-deacylating enzyme system that is active at (Hazlewood, G. P. & Dawson, R. M. C.) 49–53
Tenebrio molitor, *see* Mealworm, yellow
Testis, mouse, properties of the lactate dehydrogenase isoenzyme from (Blanco, A., Burgos, C., Gerez de Burgos, N. M. & Montamat, E. E.) 165–172
2,2,4,4-Tetrachloro-1,3-dithiacyclobutanone(dithiophosgene), inactivation by, and 2,2-dichloro-1,3-dithiacyclobutanone of ox spleen cathepsin D (Rakitzis, E. T. & Malliopoulou, T. B.) 737–739
Toxins, three, isolation, properties and amino acid sequences of, from sea-snake (*Aipysurus laevis*) venom (Maeda, N. & Tamiya, N.) 79–87

Transfer ribonucleic acid, *see* Ribonucleic acid, transfer
Transferrin, ovo-, chicken, formation of monoferric forms of, in the presence of iron-chelating agents (Donovan, J. W., Beardslee, R. A. & Ross, K. D.) 631–639
Transpeptidation, acyl and amino intermediates formed during, catalysed by pig pepsin (Wang, T. T. & Hofmann, T.) 691–699
Transpeptidation, effects of secondary binding by activator and inhibitor peptides on covalent intermediates formed during, catalysed by pig pepsin (Wang, T.-T. & Hofmann, T.) 701–712
Tricarboxylic acid cycle, factors affecting the activity of *Acetobacter xylinum* citrate synthase and its possible role in the regulation of (Swissa, M. & Benziman, M.) 173–179
Triterpenoid carotenoids, identification of 4,4′-diaponeurospen-4-al and 4,4′-diapolycopen-4-al as, in *Streptococcus faecium* UNH 564P (Taylor, R. F. & Davies, B. H.) 233–239
Troponin C, binding sites of rabbit white-skeletal-muscle troponin I for actin and (Syska, H., Wilkinson, J. M., Grand, R. J. A. & Perry, S. V.) 375–387
Troponin I, white-skeletal-muscle, rabbit, relationship between biological activity and primary structure of (Syska, H., Wilkinson, J. M., Grand, R. J. A. & Perry, S. V.) 375–387
Tubers, sweet-potato, effect of urea on the limited hydrolysis of amylose by β-amylase from (Patil, N. B.) 339–342
Turkey, studies by electron-paramagnetic-resonance spectroscopy and stopped-flow spectrophotometry on the mechanism of action of xanthine dehydrogenase from the liver of (Barber, M. J., Bray, R. C., Lowe, D. J. & Coughlan, M. P.) 297–307
Tyrosine, study of the supposed conversion of, into 3,4-dihydroxyphenylalanine catalysed by horseradish peroxidase (Smith, P. I. & Swan, G. A.) 403–408

Urine, human, chemical and physical properties of the glycoprotein with gastric anti-secretory activity from (Lugaro, G., Pasta, P., Casellato, M. M., Mazzola, G. & Carrea, G.) 641–646

Veillonella alcalescens, studied by electron-paramagnetic-resonance spectroscopy on the mechanism of action of xanthine dehydrogenase from (Dalton, H., Lowe, D. J., Pawlik, R. T. & Bray, R. C.) 287–295
Venom, European-sand-viper, purification and properties of kininogenin from (Bailey, G. S. & Shipolini, R. A.) 409–414
Venom, sea-snake (*Aipysurus laevis*), isolation, properties and amino acid sequences of three neurotoxins from (Maeda, N. & Tamiya, N.) 79–87
Venom, sea-snake (*Laticauda semifasciata*), preparation and activity of guanidinated and acetylated erabutoxins from (Hori, H. & Tamiya, N.) 217–222
Vipera ammodytes ammodytes, *see* European sand viper
Vipera berus, *see* Adder

Whey proteins, milk, red-deer, characterization of a homologue of cow β-lactoglobulin from (McDougall, E. I. & Stewart, J. C.) 647–655

Xanthine dehydrogenase, liver, turkey, studies by electron-paramagnetic-resonance spectroscopy and stopped-flow spectrophotometry on the mechanism of action of (Barber, M. J., Bray, R. C., Lowe, D. J. & Coughlan, M. P.) 297–307
Xanthine dehydrogenase, *Veillonella alcalescens*, studies by electron-paramagentic-resonance spectroscopy on the mechanism of action of (Dalton, H., Lowe, D. J., Pawlik, R. T. & Bray, R. C.) 287–295
Xanthophylls, triterpenoid, identification of 4,4′-diaponeurospen-4-al and 4,4′-diapolycopen-4-al as, in *Streptococcus faecium* UNH 564P (Taylor, R. F. & Davies, B. H.) 233–239

Yeast, association of phosphoglycerate kinase from (Spragg, S. P., Wilcox, J. K., Roche, J. J. & Barnett, W. A.) 423–428
Yeast, baker's (*Saccharomyces cerevisiae*), cyanogen bromide cleavage and the amino acid sequence of a peptide containing the active-site histidine residue of phosphoglycerate mutase from (Fothergill, L. A. & Hodgson, G. I.) 145–149
Yeast, baker's (*Saccharomyces cerevisiae*), derivation of the complete sequence of cysteine transfer ribonucleic acid from (Holness, N. J. & Atfield, G.) 447–454
Yeast, baker's (*Saccharomyces cerevisiae*), extraction and purification of a cysteine transfer ribonucleic acid from (Holness, N. J. & Atfield, G.) 429–435
Yeast, baker's (*Saccharomyces cerevisiae*), nucleotide sequences of the products obtained by the complete digestion by pancreatic ribonuclease and ribonuclease T_1 of cysteine transfer ribonucleic acid from (Holness, N. J. & Atfield, G.) 437–446
Yeast, baker's (*Saccharomyces cerevisiae*), partial purification and properties of alcohol dehydrogenase from, grown without added zinc ions (Dickenson, C. J. & Dickinson, F. M.) 309–319
Yellow mealworm, *see* Mealworm, yellow

Zinc, effects of the replacement by other metals of, in *Bacillus stearothermophilus* fructose diphosphate aldolase and a comparison of the mechanisms of action of class I and class II fructose diphosphate aldolases (Hill, H. A. O., Lobb, R. R., Sharp, S. L., Stokes, A. M., Harris, J. I. & Jack, R. S.) 551–560
Zinc ions, influence of complexing agents for, and other metal ions on the stability and activity of pig kidney alkaline phosphatase (Ackermann, B. P. & Ahlers, J.) 151–157
Zinc ions, partial purification and properties of alcohol dehydrogenase from baker's yeast grown in the absence of (Dickenson, C. J. & Dickinson, F. M.) 309–319